

QFT Problem Set 6 - Due Dec. 1

You should skim chapters 15 - 19, focusing on the material we've covered. As usual, *problems* are for extra credit seekers, although everyone should look at them.

1. **Book Problems** 15.3, 18.1, 19.1, *15.4*, *Appendix B.1*, *19.3*
2. **Integrating Out Charged Particles** Consider scalar QED, where the scalar has mass M . Are there any tree level Feynman diagrams for scattering of photons into photons (2-to-2)? What diagrams contribute to this process at one-loop? Estimate the leading result when the photons in the scattering process have momentum much less than the scale set by M . Write down the corresponding effective operator(s) that contribute to leading order in the Lagrangian. With M set equal to the electron mass, are you surprised that optical photons don't scatter off of each other?
3. **Kadanoff Block Spins and Renormalization Flow** Here we will explicitly derive a very simple example of renormalization flow. Consider a 1-d Ising model with partition function

$$Z = \sum_{S_i = \pm 1} \exp \left[\sum_i L(i, i+1) \right] \quad \text{where} \quad L(i, j) = K S_i S_j + \frac{1}{2} h (S_i + S_j) \quad (1)$$

where K and h are constants, and S_i are spin variables that can take values $S_i = \pm 1$. Since we sum over all possible states to compute the partition function, we sum over all 2^N possibilities for the N different S_i . You can assume that the chain of spins forms a circle (so we have periodic boundary conditions).

Note that the exponent of the partition function plays exactly the same role as the action for a QFT. When we discuss path integrals you will learn that statistical mechanics is basically just QFT with $t \rightarrow it$.

Let us now integrate out the even numbered spins, to get a new action for the odd spins. This is a concrete way to integrate out short distance modes and derive an effective theory for the long-distance modes.

- (a) Show that if we sum over the even spins, we are left with a new effective Lagrangian for the odd spins

$$\tilde{L}(i, i+2) = \frac{1}{2} h (S_i + S_{i+2}) + \log (2 \cosh [K(S_i + S_{i+2}) + h]) \quad (2)$$

Expand this to linear order in the external magnetic field h .

- (b) Show that

$$(S_i + S_j)^{2n} = 2^{2n-1} (1 + S_i S_j) \quad (3)$$

$$(S_i + S_j)^{2n+1} = 2^{2n} (S_i + S_j) \quad (4)$$

(c) Use this to write the new action \tilde{L} at linear order in h as

$$\tilde{L}(i, i+2) = K' S_i S_{i+2} + \frac{1}{2} h' (S_i + S_{i+2}) \quad (5)$$

You should find that

$$K' = \frac{1}{2} \log \cosh(2K) \quad (6)$$

$$h' = h[1 + \tanh(2K)] + \mathcal{O}(h^2) \quad (7)$$

We have derived the RG equations for the evolution of the lagrangian parameters K and h as we zoom out to larger and larger distances. Note that in this theory we were very lucky that the renormalized Lagrangian took the same form as the original Lagrangian – in general many other terms will be generated. Polchinski’s classic “Renormalization and Effective Lagrangians” (Nucl.Phys. B231 (1984) 269-295) applies this in a QFT context.